

THE HOLOCENE-PLEISTOCENE BOUNDARY ZONE IN THE LAGOON OF VENICE

Maurizio Bonardi* and Luigi Tosi*

* Consiglio Nazionale delle Ricerche, Istituto per lo Studio della
Dinamica delle Grandi Masse, San Polo 1364, 30125 Venezia (Italy).

Abstract

The results of a detailed study on the sediments representative of the boundary area from the Late-Pleistocene and the Lower-Holocene sedimentations are reported. Mineralogical, granulometric and geochemical investigations have been carried out to characterise the clay-rich samples taken along the littoral bar and representative of the continental and marine sedimentations. Particular attention was devoted to the study of the overconsolidated clay layer considered to be the last continental deposit before the Holocene (Flandrian) transgression and interpreted as a paleosol.

Introduction

The arch-shaped Lagoon of Venice, located in the north-western Adriatic Sea, covers an area of about 550 square Km, is separated from the open sea by a narrow littoral sand bar.

At present the lagoon communicates with the open sea by three openings that allow water exchange and maintain the marine character of the lagoon, while all the major rivers have been diverted outside the lagoon.

This study focuses on the mineralogical and textural aspects of the overconsolidated clay layer at the Holocene-Pleistocene boundary zone in comparison to the closely overlying and underlying clay-rich layers.

Late-Quaternary evolution of the Venetian lagoon

The paleoevolution of the Venetian basin has been characterised by three main episodes:

- the Pleistocene continental sedimentation during the Würm glaciation, when the climate varied from cold and dry to cold and very dry, and the Adriatic Sea level was 90 to 130 m lower than present and the coastline was about 300 km further South.
- the period of emergence of about 11,000 years (from 18,000 to 7,000 years B.P.) that characterised the final phase of the Pleistocene, with almost no sedimentation. During this time the climate improved and the carbonate muds were drained, desiccated, compacted and oxidised to form the

overconsolidated clay level representative of the Holocene/Pleistocene limit and locally known as *caranto*.

- the Holocene Flandrian transgression characterised by a warm and humid climate, during which the continental sediments were partly eroded, reworked and dispersed by the intense fluvial activity. While the climate improved the sea level began to rise over the northern Adriatic paleoplain and the coastline moved northwards until it reached approximately the present position, about 6,000 years B.P., and formed the thin sandy bar that delineates the primordial Venetian Lagoon. The intense fluvial activity, with *overflow* episodes, at first and the marine one later, partly eroded and reworked the hard clay layers leaving deep scars in the morphology.

This study focuses on the textural, mineralogical and chemical characteristics of the *caranto* layer in comparison with the other overlying and underlying clay-rich layers. The hard clay layer (*caranto*) shows a fairly high consistence due to the prolonged subareal exposure in cold and very dry climate that favoured the drainage and consolidation process. The effect of the climatic variation can be found in the mineralogical and chemical (Fig. 1,2) compositions and textural characteristics that are quite different from those of the clay layers above and below the *caranto*.




Fig. 1

Particle size distribution is shown as percentage of clay (<2 microns), silt (2-60 microns), and sand (> 60 microns) particle in each sample (Fig.1). The plot of this data shows that these samples are classified as silt to sandy silts.




Fig. 2

- Holocene
- Caranto
- △ Pleistocene





Fig. 3

Bonardi and Tosi, 1994, The Holocene-Pleistocene boundary zone in the Lagoon of Venice.
 In: Marabini F. Editor, Italian Contribution to the China-Italy Bilateral Seminar on Marine Geology,
 (4-6 October 1994, Qingdao, China). Istituto di Geologia Marina, CNR, Bologna, Tech. Rep. 40, 18-24.

Conclusions

This study has indicated that the chemical, mineralogical and textural characteristics of the clay sediments in the Holocene/Pleistocene boundary zone are closely related mainly to the paleoclimatic changes and, to a less extent, to the depositional environments.

A similar approach was used to investigate the boundary zone in the South Yellow Sea and in the Yangtze River Delta (China).

The results from both investigations indicate that exists a close correlation between paleoclimatic events and compositional and textural variations of the sediments

Acknowledgements

Financial support was provided by the research project: *Sistema Lagunare Veneziano, L.R. 2.5, U.O. 2.*

Bibliography

- BONARDI M. & CARBOGNIN L. (1992): *Late Quaternary sediments differentiation and characterisation by EMPA-EDS mineral assemblage study*. Electron Microscopy 92, Proc. of the 10th EUREM 92, Vol. 2, p. 603, Granada.
- BONARDI M. & TOSI L. (1994): *I sedimenti tardo-quadernari del cordone litoraneo della Laguna di Venezia: le sabbie*. Istituto Studio Dinamica Grandi Masse, Techn. Rep. 184, Venezia.
- BONARDI M. & TOSI L. (1994): *Climatic and sea level fluctuations evidenced in the mineralogical composition of Late Quaternary sediments from the Venetian Littoral (Italy)*. Proc. of the 14th International Sedimentological Congress, Recife, Brazil, E5.
- BONARDI M. & TOSI L. (1994): *Effects of Late Quaternary climatic changes on an exposed clay layer in the Lagoon of Venice (Italy)*. Proc. of the 14th International Sedimentological Congress, Recife, Brazil, D17.
- GATTO P. & PREVIATELLO P. (1974): *Significato stratigrafico, comportamento meccanico e distribuzione nella laguna di Venezia di un argilla sovraconsolidata nota come 'caranto'*. C.N.R., Istituto Studio Dinamica Grandi Masse, Techn. Rep. 70, Venezia.
- QIN Y., ZHAO Y., CHEN L. & ZAO S., (1990): *Geology of Bohai Sea*. China Ocean Press, Beijing.
- STANLEY D. J & CHEN Z., (1993): *Yangtze delta, eastern China: 1. Geometry and subsidence of Holocene depocenter*. Mar. Geol., 112: 1-11.
- TOSI L. (1993): *Caratteristiche geotecniche del sottosuolo del litorale veneziano*. C.N.R. Istituto per lo Studio della Dinamica delle Grandi Masse, Tech. Rep., 171, Venezia.
- TOSI L. (1994): *Rapporto e prime interpretazioni sulle analisi paleontologiche condotte su campioni tardo-quadernari del sottosuolo del litorale veneziano*. C.N.R. Istituto per lo Studio della Dinamica delle Grandi Masse, Tech. Rep., 182, Venezia.
- VISHER G. S. (1969): *Grain size distributions and deposition processes*. J. Sediment. Petrol. 39, 1074-1106.